

## Instructions For Dr. Anderson's Cataract Surgery Patients After Surgery

Continue to Use **ALL** of the following medications as directed:

1. **OCUFLOX**  
Antibiotic  
(ofloxacin)

4 x per day  
in operated eye  
for 2 weeks  
and then stop



2. **PRED-FORTE (OPHTHOTATE)**  
Anti-Inflammatory  
**AND** (prednisolone acetate)

4 x per day in operated eye  
for 2 weeks  
3 x per day for the next week  
2 x per day for the next week  
1 x per day for the next week  
and then stop



3. **VOLTAREN OPHTHA**  
Anti-Inflammatory  
**AND** (diclofenac sodium)

4 x per day  
in operated eye  
for 2 weeks  
and then stop



Be sure to wait 3 minutes between the instillation of each drop so as not wash the first drop out with the second drop.

Just before bed, apply shield (point towards nose) with scotch tape or paper tape for 1 week. The shield is to keep your hands out of your eyes while you are sleeping.

### Additional Instructions

If you experience any of the following symptoms please contact Dr. Anderson:

- severe pain
- feel sick or vomit
- notice your vision worsening instead of improving from day to day
- see flashes of light or a dramatically increased number of "floaters" (the black specks that sometimes appear in your field of vision)
- have redness that doesn't go away with the eye drops that you were given

**Please call Dr. Anderson's office at 604-530-6838**  
**After office hours, Dr. Anderson's home number is 604-881-7045**

All of Dr. Anderson's Patient Instructions can be viewed on the Internet by visiting  
<http://www.prk.com/eye-surgeons/anderson>